



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

manganese iron, especially well adapted for car-wheels, has long been prepared by the old charcoal process. Iron has been mined almost continuously since 1734. Most of the mines are located at or near the boundary of the areas of Hudson schist with those of the Stockbridge dolomite, both of which are Cambro-Ordovician sedimentaries. The more important of the exploited ore beds form a nearly continuous series encircling the base of Mt. Washington. The author believes that the ore was derived from pyrite in the Berkshire schists of the adjacent elevated territory, and that the ore bodies have been formed by the replacement of Berkshire schist and Stockbridge dolomite.

C. W. W.

Lead and Zinc Deposits of Virginia. By THOMAS LEONARD WATSON. (Geological Survey of Virginia, Geological Series, Bulletin No. 1. Pp. 156, 14 plates. Published by the Board of Agriculture and Immigration, Richmond, Va., 1905.)

Galenite and sphalerite are associated in all the mining districts, generally as replacement deposits in limestone breccia near faults, and on anticlinal axes. The sulphide ores show no secondary enrichment. The most interesting feature of this region is the secondary oxidized ore which occurs in depressions of the weathered surface of the limestone beneath several feet of residual clay. This ore consists of predominant calamine, associated with smithsonite, and cerrusite, and toward the bottom there is generally some galenite. Commercial ores are limited to the Shenandoah Limestone.

C. W. W.

The Paragenesis of the Minerals in the Glaucophane-Bearing Rocks of California. By JAMES PERRIN SMITH. (Proceedings of the American Philosophical Society, Vol. XLV, 1906, pp. 183-242.)

This paper, and the one by H. S. Washington published last year, make a nearly complete study of the glaucophane schists and related rocks. Professor Smith shows that the glaucophane rocks of the Coast Ranges have been derived from siliceous fragmental sediments, deposits of organic silica, acid arkoses, medium-basic clay shales, basic tuffs, syenites, diorites, diabases, gabbros, and probably pyroxenites. The origin may be determined by study of the chemical composition. Metamorphism has consisted merely in recrystallization, no material has been added or taken away, except that the water which once existed in the pore spaces has been included as water of crystallization. The paper includes thirty-two chemical analyses, and petrographic descriptions of the minerals and rocks.

C. W. W.